California Fish and Game 86(4):263-267 2000

INDEX TO VOLUME 86 (2000)

AUTHORS

Amador-Silva, E.: see Balart, Castro-Aguirre, and Amador-Silva.
Anonymous: Cantara spill bibliography, 101-114.
Antonio, D.B., C. Swanson, J.J. Cech Jr., R.C. Mager, S. Doroshov, and R.P. Hedrick: Prevalence of *Mycobacterium* in wild and captive delta smelt, 233-243.
Balart, E.F., J.L. Castro-Aguirre, and E. Amador-Silva: The first eastern Pacific report of the sharptail mola, *Mola lanceolata* (Tetraodontiformes: Molidae), 156-158.
Banks, J.D.: see Jones, Levine, and Banks.
Barry, J.P. and N. Maher: Observation of the prickly shark, *Echinorhinus cookei*, from the oxygen minimum zone in Santa Barbara Basin, California, 213-215.
Bleich, V.C.: see Schaefer, Torres, and Bleich.
California Bird Records Committee: Annotated list of the birds of California, 186-207.
Castro-Aguirre, J.L.: see Balart, Castro-Aguirre, and Amador-Silva.
Cech, J.J. Jr.: see Antonio and 5 others.
Corcoran, D.M.: see Crain and Corcoran.
Crain, P.K. and D.M. Corcoran: Age and growth of tui chub in Eagle Lake, California, 149-155.
Davis, L.H.: see Moyle and Davis.
Doroshov, S.: see Antonio and 5 others.
Fountain, M.C.: see Nielsen, Heine, Gan, and Fountain.
Fox, L.M., P.R. Krausman, M.L. Morrison, and R.M. Kattnig: Water and nutrient content of forage in Sonoran pronghorn habitat, Arizona, 216-232.
Fox, L.M., P.R. Krausman, M.L. Morrison, and T.H. Noon: Mineral content of Sonoran pronghorn forage, 159-174.
Gan, C.A.: see Nielsen, Heine, Gan, and Fountain.
Garrison, B.A.: Symposium overview: Damage to and recovery of aquatic and riparian ecosystems in the upper Sacramento River from the Cantara chemical spill, 1-3.
Hankin, D.G. and D. McCanne: Estimating the number of fish and crayfish killed and the proportions of wild and hatchery rainbow trout in the Cantara spill, 4-20.
Hedrick, R.P.: see Antonio and 5 others.
Heine, E.L.: see Nielsen, Heine, Gan, and Fountain.
Jones, K.C., K.F. Levine, and J.D. Banks: DNA-based genetic markers in black-tailed and mule deer for forensic applications, 115-126.
Kattnig, R.M.: see Fox, Krausman, Morrison, and Kattnig.
Krausman, P.R.: see Fox, Krausman, Morrison, and Kattnig; Fox, Krausman, Morrison, and Noon.
Levine, K.F.: see Jones, Levine, and Banks.
Lindstrand, L. III: Discovery of Shasta salamanders in atypical habitat, 259-261.
Lovich, J.E.: see Nicolai and Lovich.
Luke, C. and D. Sterner: Possible effects of the Cantara spill on amphibian populations of the upper Sacramento River, 41-60.
Luke, C. and D. Sterner: Possible impacts of the Cantara spill on reptile populations along the upper Sacramento River, 61-71.
Mager, R.C.: see Antonio and 5 others.
Maher, N.: see Barry and Maher.

Martz, C.P.: see Sheehy and 5 others.
McCanne, D.: see Hankin and McCanne.
Miller, J.W.: see Sheehy and 5 others.
Milton, J.P.: see Sheehy and 5 others.
Morrison, M.L.: see Fox, Krausman, Morrison, and Kattnig; Fox, Krausman, Morrison, and Noon.
Moyle, P.B. and L.H. Davis: A list of freshwater, anadromous, and euryhaline fishes of California, 244-258.
Nicolai, N.C. and J.E. Lovich: Preliminary observations of the behavior of male, flat-tailed horned lizards before and after an off-highway vehicle race in California, 208-212.
Nielsen, J.L., E.L. Heine, C.A. Gan, and M.C. Fountain: Molecular analysis of population genetic structure and recolonization of rainbow trout following the Cantara spill, 21-40.
Noon, T.H.: see Fox, Krausman, Morrison, and Noon.
O'Brien, J.W. and C.F. Valle: Food habits of California corbina in southern California, 136-148.
Richards, W.J.: see Vásquez-Yeomans, Valdéz-Moreno, and Richards.
Schaefer, R.J., S.G. Torres, and V.C. Bleich: Survivorship and cause-specific mortality in sympatric populations of mountain sheep and mule deer, 127-135.
Sheehy, D.J., C.P. Martz, M.C. Stopher, S.M. Turek, J.W. Miller, and J.P. Milton: Restoration planning for the Cantara metam sodium spill: A group multiattribute decision analysis approach, 72-86.
Sterner, D.: see Luke and Sterner.
Stopher. M.C.: Hindsight analysis for the Cantara spill Natural Resource Damage Assessment, 87-100.
Stopher, M.C.: see Sheehy and 5 others.
Swanson, C.: see Antonio and 5 others.
Torres, S.G.: see Schaefer, Torres, and Bleich.
Turek, S.M.: see Sheehy and 5 others.
Valdéz-Moreno, M.: see Vásquez-Yeomans, Valdéz-Moreno, and Richards.
Valle, C.F.: see O'Brien and Valle.
Vásquez-Yeomans, L., M. Valdéz-Moreno, and W.J. Richards: Description of laboratory-reared larvae of California grunion and comparisons with larvae of topsmelt and jacksmelt, 175-185.

SUBJECT

Age, tui chub: in Eagle Lake, 149-155.
Amphibians: Cantara spill effects, 41-60.
Assessment: natural resource damage, 87-100.
Behavior:
lizards, flat-tailed horned, 208-212.
shark, prickly: swimming, 213-215.
Bibliography, Cantara spill, 101-114.
Birds: annotated California list, 186-207.
Cabeza Prieta National Wildlife Refuge: Sonoran pronghorn forage, 159-174, 216-232.
Cantara metam sodium spill, 1-114.
Chub, tui: age and growth in Eagle Lake, California, 149-155.
Clams, bean: food for California corbina, 136-148.
Corbina, California: food habits, 136-148.

Crab, sand: food for California corbina, 136-148.
Crayfish, signal: number killed in the Cantara spill, 4-20.
Damage assessment, natural resources: upper Sacramento River, 87-100.
Deer,
  black-tailed: DNA-based genetic markers, 115-126.
  mule: DNA-based genetic markers, 115-126.
  mule: survivorship and cause-specific mortality, 127-135.
Development, larval: California grunion, topsmelt, and jacksmelt, 175-185.
DNA-based genetic markers: black-tailed and mule deer, 115-126.
DNA, mitochondrial: rainbow trout in upper Sacramento River, 21-40.
Eagle Lake: age and growth of tui chub, 149-155.
Ecosystems, aquatic and riparian: damage and recovery, 1-3.
Ensatina: effects of toxic spill in upper Sacramento River, 41-60.
Fishes list, California: freshwater, anadromous, and euryhaline, 244-258.
Food habits: California corbina, 136-148.
Forage, Sonoran pronghorn:
  mineral content, 159-174.
  water and nutrient content, 216-232.
Forensic applications: DNA-based genetic markers in black-tailed and mule deer, 115-126.
Genetic structure, population: rainbow trout, 21-40.
Growth, tui chub: Eagle Lake, California, 149-155.
Grunion, California: larval development, 175-185.
Habitat, atypical: Shasta salamander, 259-261.
Jacksmelt: comparison of larval development with California grunion, 175-185.
Larvae: comparison of California grunion with topsmelt and jacksmelt, 175-185.
Lion, mountain: predation on mountain sheep and mule deer, 128, 130-133.
List:
  California birds, 186-207.
  California freshwater, anadromous, and euryhaline fishes, 244-258.
Lizards, flat-tailed horned: behavior before and after an off-highway vehicle race, 208-212.
Microsatellites:
  deer, black-tailed and mule, 115-126.
  rainbow trout, 21-40.
Mola, sharptail: first eastern Pacific report, 156-158.
Mortality, cause-specific: in sympatric populations of mountain sheep and mule deer, 127-135.
Movements, flat-tailed horned lizards: effects of off-highway vehicle race, 208-212.
Mycobacterium: prevalence in delta smelt, 233-243.
Nutrients: in Sonoran pronghorn forage, 216-232.
Pacific Ocean, eastern: range extension of the sharptail mola, 156-158.
Pronghorn, Sonoran:
  mineral content of forage, 159-174.
  water and nutrient content of forage, 216-232.
Race, off-highway vehicle: effect on behavior of male, flat-tailed horned lizards, 208-212.
Range extension: sharptail mola in the eastern Pacific, 156-158.
Reptiles: Cantara spill impacts, 61-71.
Resources, natural: damage assessment, 87-100.
Restoration planning: Cantara spill, 72-86.
Sacramento River, upper: Cantara metam sodium spill, 1-114.

Salamander,
Pacific giant: recovery after the Cantara spill, 41-60.
Shasta: in atypical habitat, 259-261.
Santa Barbara Basin: prickly shark from the oxygen minimum zone, 213-215.
Shark, prickly: from the oxygen minimum zone in Santa Barbara Basin, 213-215.
Sheep, mountain: survivorship and cause-specific mortality, 127-135.
Smelt, delta: prevalence of *Mycobacterium*, 233-243.
Spill, Cantara chemical:
amphibian population effects, 41-60.
bibliography, 101-114.
fish and crayfish killed, 4-20.
Natural Resource Damage Assessment, 87-100.
overview, 1-3.
rainbow trout:
population genetic structure, 21-40.
wild and hatchery proportions, 4-20.
reptile population impacts, 61-71.
restoration planning, 72-86.
Survivorship: in sympatric populations of mountain sheep and mule deer, 127-135.
Topsmelt: comparison of larval development with California grunion, 175-185.
Trout, rainbow:
genetic structure, 21-40.
mitochondrial DNA, 21-40.
recolonization of a stream, 21-40.
wild and hatchery proportion in the Cantara spill fish kill, 4-20.

SCIENTIFIC NAMES

Aeromonas sp., 237.
Ambrosia deltoidea, 168, 223, 224.
Ambrosia dumosa, 168, 223, 224.
Ambystoma macrodactylum, 53.
Amphistichus argenteus, 145.
Aneides flavipunctatus, 50, 52.
Antilocapra americana americana, 159, 216.
Antilocapra americana mexicana, 159, 216.
Antilocapra americana oregona, 159, 216.
Antilocapra americana peninsularis, 160, 216.
Antilocapra americana sonoriensis, 159-174, 216-232.
Artemia sp., 176, 235.
Ascaphus truei, 50.
Atherinops affinis, 175-185.
Atherinops californiensis, 175-185.
Beloperone californica, 168, 223, 224.
Bufo boreas, 50.
Callianassa spp., 139.
Callisaurus draconoides, 211.
Catostomus occidentalis, 7.
Catostomus tahoensis, 150.
Charina bottae, 62.
Chasmistes cujus, 154.
Clemmys marmorata, 61, 62.
Cnemidophorus tigris, 62, 211.
Coluber constrictor, 62, 68.
Contia tenuis, 62.
Cottus gulosus, 4, 7.
Crotalus viridis, 62.
Diadophis punctatus, 62.
Dicamptodon tenebrosus, 41-60.
Donax spp., 139.
Echinorhinus brucus, 213.
Echinorhinus cookei, 213-215.
Elgaria coerulea, 62.
Elgaria multicarinata, 62.
Emerita analoga, 136.
Encelia farinosa, 168, 223, 224.
Ensatina eschscholtzii, 41-60.
Escherichia coli, 115, 118.
Eumeces skiltonianus, 62.

Euphorbia spp., 168, 223, 224.
Gila bicolor, 149-155.
Hydromantes shastae, 259-261.
Hyla regilla, 50.
Hypomesus transpacificus, 233-243.
Hyptis emoryi, 168, 223, 224.
Janusia gracilis, 168, 223, 224.
Jatropha cuneata, 168, 223, 224.
Krameria spp., 168, 223, 224.
Lampropeltis getula, 62.
Lampropeltis zonata, 62.
Larrea tridentata, 168, 223, 224.
Leuresthes tenuis, 175-185.
Lutra canadensis, 59.
Masticophis lateralis, 62.
Masturus lanceolata, 156.
Masturus oxyuropterus, 156.
Menticirrhus americanus, 145.
Menticirrhus littoralis, 145.
Menticirrhus ophicephalus, 144.
Menticirrhus undulatus, 136-148.
Micropterus dolomieu, 7, 96.
Micropterus punctulatus, 7, 96.
Morone saxatilis, 234.
Mycobacterium chelonae, 233, 237, 238, 240, 241.
Mycobacterium marinum, 238.
Mycobacterium sp., 233-243.
Myliobatis californica, 145.
Mylopharodon conocephalus, 7.
Odocoileus hemionus, 127-135.
Odocoileus hemionus columbianus, 115-126, 170.
Odocoileus hemionus eremicus, 227.
Odocoileus hemionus hemionus, 115-126.
Olneya tesota, 168, 223, 224.
Oncorhynchus mykiss, 4-20, 21-40, 96.
Oncorhynchus mykiss aquabonita, 30.
Oncorhynchus mykiss aquilarium, 30, 149.
Oncorhynchus nerka, 26.
Oncorhynchus tshawytscha, 26.
Opuntia fulgida, 168, 223, 224.
Ovis canadensis, 127-135.
Ovis canadensis californiana, 133.
Ovis canadensis cremnobates, 133.
Pagurus spp., 139.
Pascifastacus leniusculus, 7, 62.
Phrynosoma mcallii, 208-212.
Pituophis melanoleucus, 62.
Prosopis juliflora, 168, 223, 224.
Ptychocheilus grandis, 7.
Puma concolor, 128.
Rana aurora, 53.
Rana boylii, 50.
Rana cascadae, 53.
Rana catesbeiana, 50.
Rhinichthys osculus, 150.
Richardsonius egregius, 150.
Salmo salar, 26.
Salmo trutta, 7, 96.
Salvelinus fontinalis, 7, 96.
Sapium bioculare, 168, 223, 224.
Sceloporus graciosus, 62.
Sceloporus occidentalis, 61, 62, 70.
Sphaeralcea spp., 168, 223, 224.
Solen rosaceus, 139.
Syngnathus spp., 139.
Tagelus californianus, 139.
Taricha granulosus, 50.
Tetraselmis suecica, 176.
Thamnophis atratus, 62.
Thamnophis elegans, 62, 68.
Thamnophis sirtalis, 62.
Umbrina roncader, 145.
Urolophus halleri, 145.
Uta stansburiana, 211.